

Isospin Symmetry Breaking in the Quark Condensates, $\langle \bar{d}d \rangle \neq \langle \bar{u}u \rangle$, as Estimated from QCD Sum Rules

W-Y. P. Hwang[†] and Kwei-Chou Yang[‡]

[†]*Department of Physics, National Taiwan University, Taipei, Taiwan 10764, R.O.C.*

[‡]*Institute of Physics, Academia Sinica, Taipei, Taiwan 115, R.O.C.*

Abstract

Isospin symmetry breaking in the quark condensates, $\langle \bar{d}d \rangle \neq \langle \bar{u}u \rangle$, is a fundamental parameter in both the QCD sum rule studies and the chiral perturbation theory. In this article, we apply the QCD sum rule method to treat the hyperon mass difference $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}$ in order to obtain a reasonable estimate on such isospin symmetry breaking parameter γ defined as $(\langle \bar{d}d \rangle - \langle \bar{u}u \rangle)/\langle \bar{u}u \rangle$. Note that the electromagnetic contributions to the particular mass difference are expected to cancel almost completely. Using the light-quark mass difference $m_d - m_u = 4 \pm 1$ MeV and the experimental data $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+} = 5.62 \pm 0.13$ MeV, we obtain $\gamma = -0.011 \pm 0.001$, a value which is slightly larger than the commonly adopted value and is of smaller uncertainty.

PACS number(s): 11.30.Hv; 12.38.Lg; 11.55Hx; 14.20.-c

Although quantum chromodynamics (QCD) has been accepted as a theory for describing strong interactions among quarks and gluons, it is highly nonperturbative at the confinement scale. In particular, it is expected that the QCD ground state, or the QCD vacuum, has nonzero quark condensate $\langle \bar{q}q \rangle$, gluon condensate $\langle G^2 \rangle$, and many other condensates of higher order. Making use of the method QCD sum rules[1], it has become possible to study hadron physics in terms of the basic parameters of QCD, such as current quark masses and non-perturbative condensates. In a systematic study of isospin symmetry breaking effects, both the quark mass difference, $m_d - m_u$, and the difference in condensates, $\gamma \equiv (\langle \bar{d}d \rangle / \langle \bar{u}u \rangle) - 1$ are the most fundamental parameters. while the quark mass difference $m_d - m_u$ can be estimated with reasonable confidence as $m_d - m_u \approx 3 - 5$ MeV[2, 3], γ is poorly known. Theoretical estimates on γ have been obtained in the chiral perturbation theory[4], in the Nambu-Jona-Lasinio model[5, 6], and in QCD sum rules[7, 8, 9, 10, 11]; the value has a fairly large range, $\gamma = -(2 \sim 10) \times 10^{-3}$. In the QCD sum rule approach, the error in estimating the value of γ comes from the poorly-known electromagnetic contributions[8] and instanton effects[11].

In this paper, we wish to use the QCD sum rule method to obtain the isospin symmetry breaking parameter γ from the hyperon mass difference $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}$, a combination that the electromagnetic contributions are expected to cancel almost exactly. The cancelation may be understood as follows. In the simple (constituent) quark model [12, 13, 2, 3], the electromagnetic contributions can be separated into a contribution from photon exchange between two of the three quarks and a contribution from the quark self-energy type:

$$m^\gamma = \alpha \langle 1/r \rangle (Q_1 Q_2 + Q_2 Q_3 + Q_3 Q_1) + C(Q_1^2 + Q_2^2 + Q_3^2), \quad (1)$$

where Q_1 , Q_2 , and Q_3 are the quark charges and $\langle 1/r \rangle$ is the (effective) average inverse distance of two quarks in a baryon. In the isospin symmetric limit, the radii of Σ^+ , Σ^- , and Σ^0 are identical so that

$$\begin{aligned} \langle 1/r \rangle_{\Sigma^+} &\approx \langle 1/r \rangle_{\Sigma^-} \approx \langle 1/r \rangle_{\Sigma^0}, \\ C^{\Sigma^+} &\approx C^{\Sigma^-} \approx C^{\Sigma^0}. \end{aligned} \quad (2)$$

Here the equality would take hold should isospin symmetry be exact. We therefore obtain

$$\frac{1}{2}(m_{\Sigma^-}^{\gamma} + m_{\Sigma^0}^{\gamma}) - m_{\Sigma^+}^{\gamma} = 0. \quad (3)$$

We note that similar arguments may be constructed for other models of hadrons. Using the quark model estimates and the experimental information from electron scattering on the protons and neutrons (extrapolated to the other members of the baryon octet), Gasser and Leutwyler[3] have reported $\frac{1}{2}(m_{\Sigma^-}^{\gamma} + m_{\Sigma^0}^{\gamma}) - m_{\Sigma^+}^{\gamma} = -0.32 \pm 0.26$ MeV, which is consistent with the above observation. Accordingly, the experimentally observed value, $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+} = 5.62 \pm 0.13$ MeV, may serve to determine $m_d - m_u$ and γ in an unambiguous manner. As some authors have indicated that instanton contributions are absent in the chirally even baryon sum rules[11] but may be of significance in the chirally odd one, we choose to employ solely the chirally even sum rules.

We turn our attention to the derivation of QCD sum rules. Following the standard procedure, we consider the two-point Green's function

$$\Pi(p) = i \int d^4x \, e^{ipx} \langle 0 | T(\eta(x) \bar{\eta}(0)) | 0 \rangle, \quad (4)$$

where the composite field operators are specified by

$$\begin{aligned} \eta_{\Sigma^+} &= \epsilon^{abc} (u^{aT} C \gamma_{\mu} u^b) \gamma_5 \gamma^{\mu} s^c, \\ \eta_{\Sigma^0} &= \epsilon^{abc} \frac{1}{\sqrt{2}} [(u^{aT} C \gamma_{\mu} d^b) \gamma_5 \gamma^{\mu} s^c + (d^{aT} C \gamma_{\mu} u^b) \gamma_5 \gamma^{\mu} s^c], \\ \eta_{\Sigma^-} &= \epsilon^{abc} (d^{aT} C \gamma_{\mu} d^b) \gamma_5 \gamma^{\mu} s^c, \end{aligned} \quad (5)$$

with a , b , and c color indices and $C = -C^T$ the charge conjugate operator. At the hadronic level, we write the Green's function via dispersion relation as

$$\Pi(p) = \frac{1}{\pi} \int_0^{\infty} \frac{\text{Im } \Pi(p')}{p'^2 - p^2 - i\varepsilon} dp'^2, \quad (6)$$

where

$$\text{Im } \Pi(p') = \pi \sum_X \delta(p'^2 - M_X^2) \langle 0 | \eta(0) | X \rangle \langle X | \bar{\eta}(0) | 0 \rangle, \quad (7)$$

and

$$\langle 0|\eta(0)|\Sigma\rangle = \lambda_\Sigma v(p), \quad (8)$$

with $v(p)$ the Dirac spinor normalized as $\bar{v}(p)v(p)=2m_\Sigma$. At the quark level, we write, using the Σ^+ case as our example,

$$\langle 0|T(\eta(x)\bar{\eta}(0))|0\rangle = 2i\epsilon^{abc}\epsilon^{a'b'c'}Tr[S_u^{aa'}(x)\gamma_\nu C(S_u^{bb'}(x))^T C\gamma_\mu]\gamma_5\gamma^\mu S_d^{cc'}(x)\gamma^\nu\gamma_5. \quad (9)$$

Here we have used the definition,

$$iS^{ab}(x) \equiv \langle 0|T[q^a(x)\bar{q}^b(0)]|0\rangle,$$

as in Ref. [8].

$\Pi(p)$ has two invariant structures:

$$\Pi(p) = \Pi_1(p^2) + \Pi_{\not{p}}(p^2) \not{p}, \quad (10)$$

where Π_1 and $\Pi_{\not{p}}$ are two invariant functions. We name Π_1 as the chirally odd sum rule since chiral-odd operators contribute dominantly, and $\Pi_{\not{p}}$ as the chirally even sum rule. The derivation of the relevant QCD sum rules is similar to what we have reported in our earlier papers [8, 14]. Here we record only the final results on the $\Pi_{\not{p}}$ (chirally even) Σ mass sum rules:

$$\begin{aligned} \beta_{\Sigma^+}^2 e^{-(M_{\Sigma^+}^2/M^2)} = & \frac{M^6}{8}L^{-4/9}E_2^{\Sigma^+} + \frac{bM^2}{32}L^{-4/9}E_0^{\Sigma^+} + \frac{a_u^2}{6}L^{4/9} - \frac{a_u^2 m_0^2}{24M^2}L^{-2/27} \\ & - \frac{m_s a_s M^2}{4}L^{-4/9}E_0^{\Sigma^+} - \frac{m_s a_s m_0'^2}{24}L^{-26/27} - \frac{m_u a_u m_0^2}{12}L^{-26/27}, \end{aligned} \quad (11)$$

$$\begin{aligned} \beta_{\Sigma^0}^2 e^{-(M_{\Sigma^0}^2/M^2)} = & \frac{M^6}{8}L^{-4/9}E_2^{\Sigma^0} + \frac{bM^2}{32}L^{-4/9}E_0^{\Sigma^0} + \frac{a_u a_d}{6}L^{4/9} - \frac{a_u a_d m_0^2}{24M^2}L^{-2/27} \\ & - \frac{[m_s a_s - (a_u - a_d)(m_d - m_u)]M^2}{4}L^{-4/9}E_0^{\Sigma^0} - \frac{m_s a_s m_0'^2}{24}L^{-26/27} \\ & - \frac{[a_d(3m_u - m_d) + a_u(3m_d - m_u)]m_0^2}{48}L^{-26/27}, \end{aligned} \quad (12)$$

and

$$\beta_{\Sigma^-}^2 e^{-(M_{\Sigma^-}^2/M^2)} = \frac{M^6}{8}L^{-4/9}E_2^{\Sigma^-} + \frac{bM^2}{32}L^{-4/9}E_0^{\Sigma^-} + \frac{a_d^2}{6}L^{4/9} - \frac{a_d^2 m_0^2}{24M^2}L^{-2/27}$$

$$-\frac{m_s a_s M^2}{4} L^{-4/9} E_0^{\Sigma^-} - \frac{m_s a_s m_0'^2}{24} L^{-26/27} - \frac{m_d a_d m_0'^2}{12} L^{-26/27}, \quad (13)$$

where we have adopted the definitions:

$$\begin{aligned} a_q &\equiv -(2\pi)^2 \langle \bar{q} q \rangle, \\ a_q m_0'^2 &\equiv (2\pi)^2 \langle g_s \bar{q} \sigma \cdot G q \rangle, \quad q = u \text{ or } d, \\ b &\equiv \langle g_s^2 G^2 \rangle, \\ a_s &\equiv -(2\pi)^2 \langle \bar{s} s \rangle, \\ a_s m_0'^2 &\equiv (2\pi)^2 \langle g_s \bar{s} \sigma \cdot G s \rangle, \\ \beta_\Sigma^2 &\equiv (2\pi)^4 \frac{\lambda_\Sigma^2}{4}, \\ E_n^{(m)} &\equiv 1 - e^{-x_{(m)}} (1 + x_{(m)} + \dots + \frac{1}{n!} x_{(m)}^n), \quad \text{with } x_{(m)} = W_{(m)}^2 / M^2, \\ L &\equiv \frac{\ln(M^2 / \Lambda^2)}{\ln(\mu^2 / \Lambda^2)}. \end{aligned} \quad (14)$$

In Eqs. (11-13), M^2 is the Borel mass squared, as arising from the Borel transform of the sum rules,

$$B[\Pi_{1(p)}(p^2)] \equiv \lim_{\substack{n \rightarrow \infty \\ -p^2 \rightarrow \infty \\ \frac{-p^2}{nM^2} \text{ fixed}}} \frac{1}{n!} (-p^2)^{n+1} \left(\frac{d}{dp^2} \right)^n \Pi_{1(p)}(p^2). \quad (15)$$

In what follows, we take the value $\langle \bar{s} s \rangle = 0.8 \times (\langle \bar{u} u \rangle + \langle \bar{d} d \rangle) / 2$. We also use the relation

$$\langle g_s \bar{s} \sigma \cdot G s \rangle / \langle g_s \bar{u} \sigma \cdot G u \rangle \simeq \langle \bar{s} s \rangle / \langle \bar{u} u \rangle,$$

which indicates $m_0'^2 \simeq m_0^2$, a relation which has been examined in some detail in octet baryons [7]. The input parameters in the numerical analysis are taken to be $\bar{a}_q = (a_u + a_d) / 2 = 0.545 \text{ GeV}^3$, $a_s = 0.8 \bar{a}_q$, $b = 0.474 \text{ GeV}^4$, $m_0^2 = 0.6\text{--}0.8 \text{ GeV}^2$, $\bar{m}_q = (m_u + m_d) / 2 = 7 \text{ MeV}$, $m_d - m_u = 3\text{--}5 \text{ MeV}$, and $m_s = 150\text{--}200 \text{ MeV}$. These values correspond to the QCD scale parameters $\Lambda = 0.1 \text{ GeV}$, and the normalization point $\mu = 0.5 \text{ GeV}$. Applying the operator $M^4 \partial \ell n / \partial M^2$ to both sides of Eqs. (11-13) (to get rid of the dependence on the parameter β_Σ^2), we deduce the QCD

sum rule on the mass difference $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}$. We fit the two sides of this sum rule by minimizing

$$\delta(M^2) = \sum_{\substack{M^2=1.0+0.005i \\ 0 \leq i \leq 60}} |LHS - RHS|$$

within the Borel region $1.0 \text{ GeV}^2 \leq M^2 \leq 1.3 \text{ GeV}^2$, where the *LHS* (the left-hand side of the sum rule) is the experimental value, while the *RHS* (the right-hand side of the sum rule) is the operator-product-expansion results at the quark-gluon level. In the numerical analysis, we adopt $W^{\Sigma^+}=3.47 \text{ GeV}^2$ [14] and the experimental data $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}=5.62 \pm 0.13 \text{ MeV}$. Our results are $(W_{\Sigma^0}^2 + W_{\Sigma^-}^2)/2=3.483 \pm 0.002 \text{ GeV}^2$, $W_{\Sigma^-}^2 - W_{\Sigma^0}^2=0.0030 \pm 0.0004 \text{ GeV}^2$, and $\gamma = -0.011 \pm 0.001$. Theoretical uncertainties in our determination of the isospin symmetry breaking parameter γ come mainly from the uncertainty in the value of $m_d - m_u$ and the experimental error in $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}$.

In Fig. 1, we plot the *LHS* and *RHS* as a function of the Borel mass squared, M^2 . The solid curve is the *LHS* assuming the value $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}=5.62 \text{ MeV}$. The dotted curve is the *RHS* with the optimized values, $\gamma = -0.0114$, $W_{\Sigma^0}^2=3.4816 \text{ GeV}^2$, and $W_{\Sigma^-}^2=3.4846 \text{ GeV}^2$, where we have used, as the input, $m_0 = 0.8 \text{ GeV}^2$, $m_s = 175 \text{ MeV}$, $m_d=8.9 \text{ MeV}$, and $m_u=5.1 \text{ MeV}$.

Following the same procedure as outlined above and using the present value for γ , we have repeated the analysis of the sum rule for the neutron-proton mass difference within the Borel region $0.9 \text{ GeV}^2 \leq M^2 \leq 1.1 \text{ GeV}^2$ [8, 14]. The result is illustrated in Fig. 2, in which the solid curve is the *LHS* assuming the value $m_N^{QCD} - m_P^{QCD}=2.693 \text{ MeV}$ and the dotted curve is the *RHS* with the optimized values, $\gamma = -0.0102$, $W_N^2=2.2506 \text{ GeV}^2$ (neutron), $W_P^2=2.25 \text{ GeV}^2$ (proton), $m_d=8.9 \text{ MeV}$ and $m_u=5.1 \text{ MeV}$. Such value implies that the electromagnetic contribution to the neutron-proton mass difference $m_N^\gamma - m_P^\gamma$ is about $-1.3 \sim -1.8 \text{ MeV}$, a value in reasonable agreement with the naive quark model result [3]. Note that our result on γ is a little larger than that obtained by Forkel and Nielsen[11], who obtained $\gamma = -(0.8 - 1.0) \times 10^{-2}$ making use of $m_N^\gamma - m_P^\gamma = -0.76 \pm 0.3 \text{ MeV}$ [3] (corresponding to i.e., $m_N^{QCD} - m_P^{QCD} = 2.05 \pm 0.3 \text{ MeV}$) in the Borel region $0.8 \text{ GeV}^2 \leq M^2 \leq 1.4 \text{ GeV}^2$. Using

their value on $m_N^\gamma - m_P^\gamma$ and adopting our approach together with the same parameters as in the present paper, we obtain $\gamma = -(0.7 - 0.9) \times 10^{-2}$, a result consistent with that in Ref.[11] and also with that in our earlier paper [8]. Nevertheless, the choice $m_N^\gamma - m_P^\gamma = -0.76 \pm 0.3$ MeV can only be marginally consistent with the result obtained through the analysis of the sum rule for $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}$, which gives a better determination of γ and leads to $m_N^\gamma - m_P^\gamma$ is about $-1.3 \sim -1.8$ MeV.

In summary, we have presented a detailed analysis of the isospin symmetry breaking parameter in the quark condensate, γ , from the chirally even QCD sum rules on the mass difference $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}$, for which the instanton contributions are absent and the electromagnetic contributions are expected to cancel almost exactly. Our numerical analysis indicates that a value of γ ($= -0.011 \pm 0.001$) (slightly larger in magnitude) is preferred over the existing result, $\gamma = -(2 \sim 10) \times 10^{-3}$. Using the present value of γ and the observed neutron-proton mass difference, we have also deduced the electromagnetic contribution to the neutron-proton mass difference as in the range $-1.3 \sim -1.8$ MeV, in reasonable agreement with the quark model result.

This work was supported in part by the National Science Council of R.O.C. under Grant No. NSC87-2112-M001-048 and No. NSC87-2112-M002-031.

References

- [1] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. B**147**, 385, 448 (1979);
L. J. Reinders, H. Rubinstein, and S. Yazaki, Phys. Rep. **127**, 1 (1985).
- [2] K. Lane and S. Weinberg, Phys. Rev. Lett. **37**, 717 (1976).
- [3] J. Gasser and H. Leutwyler, Phys. Rep. **87**, 77 (1982).
- [4] J. Gasser and H. Leutwyler, Nucl. Phys. **B250**, 465 (1985).

- [5] T. Hatsuda, H. Hogaasen, and M. Prakash, Phys. Rev. C**42**, 2212 (1990).
- [6] C. Adami and G. E. Brown, Z. Phys. A **340**, 93 (1991).
- [7] V. L. Chernyak and A. R. Zhitnitsky, Phys. Rep. **112**, 173 (1984).
- [8] K.-C. Yang, W-Y. P. Hwang, E. M. Henley, and L. S. Kisslinger, Phys. Rev. D**47**, 3001 (1993).
- [9] C. Adami, E. G. Drukarev, and B. L. Ioffe, Phys. Rev. D**48**, 2304, (1993).
- [10] V. L. Eletsky and B. L. Ioffe, Phys. Rev. D**48**, 1441 (1993).
- [11] H. Forkel and M. Nielsen, Phys. Rev. D**55**, 71 (1997).
- [12] Y. Miyamoto, Progr. Theor. Phys. **35**, 175 (1966).
- [13] A. De Rújula, H. Georgi and S. L. Glashow, Phys. Rev. D**12**, 147 (1975).
- [14] W-Y. P. Hwang and K.-C. Yang, Phys. Rev. D**49**, 460 (1994).




Figure 1: $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+}$ as a function of the Borel mass squared M^2 . The solid curve is the *LHS*, the left-hand side of the sum rule, assuming $\frac{1}{2}(m_{\Sigma^-} + m_{\Sigma^0}) - m_{\Sigma^+} = 5.62$ MeV. The dotted curve is the *RHS*, obtained by making use of $m_0 = 0.8$ GeV², $m_s = 175$, $m_d = 8.9$ MeV, and $m_u = 5.1$ MeV as the input.




Figure 2: $m_N^{QCD} - m_P^{QCD}$ as a function of the Borel mass squared M^2 . The solid curve is the *LHS* assuming $m_N^{QCD} - m_P^{QCD} = 2.693$ MeV and the dotted curve is the *RHS* with the optimized values, $\gamma = -0.0102$ and $W_N^2 = 2.2506$ GeV². Here we have used $W_P^2 = 2.25$ GeV², $m_d = 8.9$ MeV and $m_u = 5.1$ MeV as the input.